

T1 Marking Scheme

Particles from the Sun

Photons from the surface of the Sun and neutrinos from its core can tell us about solar temperatures and also confirm that the Sun shines because of nuclear reactions.

Throughout this problem, take the mass of the Sun to be $M_{\odot} = 2.00 \times 10^{30}$ kg, its radius, $R_{\odot} = 7.00 \times 10^8$ m, its luminosity (radiation energy emitted per unit time), $L_{\odot} = 3.85 \times 10^{26}$ W, and the Earth-Sun distance, $d_{\odot} = 1.50 \times 10^{11}$ m.

Note:

- (i) $\int x e^{ax} dx = \left(\frac{x}{a} - \frac{1}{a^2} \right) e^{ax} + \text{constant}$
- (ii) $\int x^2 e^{ax} dx = \left(\frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right) e^{ax} + \text{constant}$
- (iii) $\int x^3 e^{ax} dx = \left(\frac{x^3}{a} - \frac{3x^2}{a^2} + \frac{6x}{a^3} - \frac{6}{a^4} \right) e^{ax} + \text{constant}$

A. Radiation from the Sun :

- (A1) Assume that the Sun radiates like a perfect blackbody. Use this fact to calculate the temperature, T_s , of the solar surface. [0.3]

Solution:

$$\text{Stefan's law: } L_{\odot} = (4\pi R_{\odot}^2)(\sigma T_s^4)$$

$$T_s = \left(\frac{L_{\odot}}{4\pi R_{\odot}^2 \sigma} \right)^{1/4} = 5.76 \times 10^3 \text{ K}$$

0.2

0.1

The spectrum of solar radiation can be approximated well by the Wien distribution law. Accordingly, the solar energy incident on any surface on the Earth per unit time per unit frequency interval, $u(\nu)$, is given by

$$u(\nu) = A \frac{R_{\odot}^2}{d_{\odot}^2} \frac{2\pi h}{c^2} \nu^3 \exp(-h\nu/k_B T_s),$$

where A is the area of the surface normal to the direction of the incident radiation.

Now, consider a solar cell which consists of a thin disc of semiconducting material of area, A , placed perpendicular to the direction of the Sun's rays.

- (A2) Using the Wien approximation, express the total power, P_m , incident on the surface of the solar cell, in terms of A , R_{\odot} , d_{\odot} , T_s and the fundamental constants c , h , k_B . [0.3]

Solution:

$$P_{\text{in}} = \int_0^{\infty} u(\nu) d\nu = \int_0^{\infty} A \frac{R_{\odot}^2}{d_{\odot}^2} \frac{2\pi h}{c^2} \nu^3 \exp(-h\nu/k_B T_s) d\nu \quad 0.1$$

Let $x = \frac{h\nu}{k_B T_s}$. Then, $\nu = \frac{k_B T_s}{h} x$ $d\nu = \frac{k_B T_s}{h} dx$.

$$P_{\text{in}} = \frac{2\pi h A R_{\odot}^2}{c^2 d_{\odot}^2} \frac{(k_B T_s)^4}{h^4} \int_0^{\infty} x^3 e^{-x} dx = \frac{2\pi k_B^4}{c^2 h^3} T_s^4 A \frac{R_{\odot}^2}{d_{\odot}^2} \cdot 6 = \frac{12\pi k_B^4}{c^2 h^3} T_s^4 A \frac{R_{\odot}^2}{d_{\odot}^2} \quad 0.2$$

Alternative Solution NOT using the Wien approximation:

$$P_{\text{in}} = \frac{L_{\odot}}{4\pi d_{\odot}^2} A = \sigma T_s^4 A \frac{R_{\odot}^2}{d_{\odot}^2} \quad (0.1)$$

$$= \left(\frac{2\pi^5 k_B^4}{15 c^2 h^3} \right) T_s^4 A \frac{R_{\odot}^2}{d_{\odot}^2} \quad (0.2)$$

σ by dimensional analysis gets 0.1 in the last step, if the entire numerical factor is not correct (the powers of all parameters mentioned in the question should be correct).

- (A3) Express the number of photons, $n_{\gamma}(\nu)$, per unit time per unit frequency interval incident on the surface of the solar cell in terms of A , R_{\odot} , d_{\odot} , T_s , ν and the fundamental constants c , h , k_B . [0.2]

Solution:

$$n_{\gamma}(\nu) = \frac{u(\nu)}{h\nu} \quad 0.1$$

$$= A \frac{R_{\odot}^2}{d_{\odot}^2} \frac{2\pi}{c^2} \nu^2 \exp(-h\nu/k_B T_s) \quad 0.1$$

The semiconducting material of the solar cell has a “band gap” of energy, E_g . We assume the following model. Every photon of energy $E \geq E_g$ excites an electron across the band gap. This electron contributes an energy, E_g , as the useful output energy, and any extra energy is dissipated as heat (not converted to useful energy).

- (A4) Define $x_g = h\nu_g/k_B T_s$ where $E_g = h\nu_g$. Express the useful output power of the cell, P_{out} , in terms of x_g , A , R_{\odot} , d_{\odot} , T_s and the fundamental constants c , h , k_B . [1.0]

Solution:

The useful power output is the useful energy quantum per photon, $E_g \equiv h\nu_g$, multiplied by

the number of photons with energy, $E \geq E_g$,

$$P_{\text{out}} = h\nu_g \int_{\nu_g}^{\infty} n_\nu(\nu) d\nu \quad 0.5$$

$$= h\nu_g A \frac{R_{\odot}^2}{d_{\odot}^2} \frac{2\pi}{c^2} \int_{\nu_g}^{\infty} \nu^2 \exp(-h\nu/k_B T_s) d\nu \quad 0.1$$

$$= k_B T_s x_g A \frac{R_{\odot}^2}{d_{\odot}^2} \frac{2\pi}{c^2} \left(\frac{k_B T_s}{h}\right)^3 \int_{x_g}^{\infty} x^2 e^{-x} dx \quad 0.4$$

$$= \frac{2\pi k_B^4}{c^2 h^3} T_s^4 A \frac{R_{\odot}^2}{d_{\odot}^2} x_g (x_g^2 + 2x_g + 2) e^{-x_g}$$

If the answer to the first part is only partially correct:

0.3 for $h\nu_g$ outside the integral sign

0.2 for the correct lower limit of the integral correct

(A5) Express the efficiency, η , of this solar cell in terms of x_g . [0.2]

Solution:

$$\text{Efficiency, } \eta = \frac{P_{\text{out}}}{P_{\text{in}}} = \frac{x_g}{6} (x_g^2 + 2x_g + 2) e^{-x_g} \quad 0.2$$

Carry over from the Alternative Solution of (A2):

$$\begin{aligned} \eta &= \frac{P_{\text{out}}}{P_{\text{in}}} = \frac{2\pi k^4}{c^2 h^3} \frac{1}{\sigma} x_g (x_g^2 + 2x_g + 2) e^{-x_g} \\ &= \frac{15}{\pi^4} x_g (x_g^2 + 2x_g + 2) e^{-x_g} = \left(\frac{90}{\pi^4}\right) \frac{x_g}{6} (x_g^2 + 2x_g + 2) e^{-x_g} \end{aligned}$$

The two answers are close, since $\frac{90}{\pi^4} = 0.92$. (0.2)

Incorrect value of η calculated correctly as the ratio of earlier calculated P_{out} and P_{in} will get credit ONLY IF it is dimensionless.

(A6) Make a qualitative sketch of η versus x_g . The values at $x_g = 0$ and $x_g \rightarrow \infty$ should be clearly shown. What is the slope of $\eta(x_g)$ at $x_g = 0$ and $x_g \rightarrow \infty$? [1.0]

Solution:


$$\eta = \frac{1}{6} (x_g^3 + 2x_g^2 + 2x_g) e^{-x_g}$$

Put limiting values, $\eta(0) = 0$ $\eta(\infty) = 0$.

Since the polynomial has all positive coefficients, it increases monotonically; the exponential function decreases monotonically. Therefore, η has only one maximum.

$$\frac{d\eta}{dx_g} = \frac{1}{6} (-x_g^3 + x_g^2 + 2x_g + 2) e^{-x_g}$$

$$\left. \frac{d\eta}{dx_g} \right|_{x_g=0} = \frac{1}{3} \quad \left. \frac{d\eta}{dx_g} \right|_{x_g \rightarrow \infty} = 0$$



Carry over from the Alternative Solution of (A2):

$$\frac{d\eta}{dx_g} = \frac{15}{\pi^4}(-x_g^3 + x_g^2 + 2x_g + 2)e^{-x_g}$$

$$\left. \frac{d\eta}{dx_g} \right|_{x_g=0} = \frac{30}{\pi^4} = 0.31 \quad \left. \frac{d\eta}{dx_g} \right|_{x_g \rightarrow \infty} = 0$$

0.2 mark EACH for the correct values and slopes at $x_g = 0$ and $x_g \rightarrow \infty$.

0.8

A smooth curve with ONE and ONLY ONE maximum.

0.2

If the expression for η is incorrect but carried forward correctly, credit may be given for slopes at 0 and ∞ .

However, the value of η has to be 0 at $x_g = 0$ and $x_g \rightarrow \infty$ since these are the only physically meaningful answers.

- (A7) Let x_0 be the value of x_g for which η is maximum. Obtain the cubic equation that gives x_0 . Estimate the value of x_0 within an accuracy of ± 0.25 . Hence calculate $\eta(x_0)$.

[1.0]

Solution:

The maximum will be for $\frac{d\eta}{dx_g} = \frac{1}{6}(-x_g^3 + x_g^2 + 2x_g + 2)e^{-x_g} = 0$

$$\Rightarrow p(x_g) \equiv x_g^3 - x_g^2 - 2x_g - 2 = 0$$

0.4

A Numerical Solution by the Bisection Method:

Now,

$$\begin{aligned} p(0) &= -2 \\ p(1) &= -4 \\ p(2) &= -2 \\ p(3) &= 10 \quad \Rightarrow \quad 2 < x_0 < 3 \\ p(2.5) &= 2.375 \quad \Rightarrow \quad 2 < x_0 < 2.5 \\ p(2.25) &= -0.171 \quad \Rightarrow \quad 2.25 < x_0 < 2.5 \end{aligned}$$

The approximate value of x_g where η is maximum is $x_0 = 2.27$.

Alternative methods leading to the same result are acceptable.

Answer in the range $1.95 \leq x_0 \leq 2.55$

0.4

Answer in the range $1.75 \leq x_0 < 1.95$ or $2.55 < x_0 \leq 2.75$

(0.3)

Answer in the range $1.50 \leq x_0 < 1.75$ or $2.75 < x_0 \leq 3.00$

(0.2)

The above 0.4 marks for the numerical answer to be given only if the numerical analysis is done for a cubic equation. Solving for an incorrect linear / quadratic or similar simpler equation does not get credit.

However, if instead of obtaining a cubic equation, the original η has been maximised by numerical means and the answer is correct, then these 0.4 marks may be given.

$$\eta(2.27) = 0.457 \text{ (two significant figures accepted, range } 0.45 - 0.47 \text{ OK)}$$

0.2

- (A8) The band gap of pure silicon is $E_g = 1.11$ eV. Calculate the efficiency, η_{Si} , of a silicon solar cell using this value. [0.2]

Solution:

$$x_g = \frac{1.11 \times 1.60 \times 10^{-19}}{1.38 \times 10^{-23} \times 5763} = 2.23$$

0.1

$$\eta_{\text{Si}} = \frac{x_g}{6} (x_g^2 + 2x_g + 2) e^{-x_g} = 0.457$$

0.1

Carry over from the Alternative Solution of (A2):

$$\eta_{\text{Si}} = \frac{15x_g}{\pi^4} (x_g^2 + 2x_g + 2) e^{-x_g} = 0.422$$

(0.1)

In the late nineteenth century, Kelvin and Helmholtz (KH) proposed a hypothesis to explain how the Sun shines. They postulated that starting as a very large cloud of matter of mass, M_\odot , and negligible density, the Sun has been shrinking continuously. The shining of the Sun would then be due to the release of gravitational energy through this slow contraction.

- (A9) Let us assume that the density of matter is uniform inside the Sun. Find the total gravitational potential energy, Ω , of the Sun at present, in terms of G , M_\odot and R_\odot . [0.3]

Solution:

The total gravitational potential energy of the Sun: $\Omega = - \int_0^{M_\odot} \frac{Gm \, dm}{r}$

0.1

For constant density, $\rho = \frac{3M_\odot}{4\pi R_\odot^3}$ $m = \frac{4}{3}\pi r^3 \rho$ $dm = 4\pi r^2 \rho dr$

$$\Omega = - \int_0^{R_\odot} G \left(\frac{4}{3}\pi r^3 \rho \right) (4\pi r^2 \rho) \frac{dr}{r} = - \frac{16\pi^2 G \rho^2 R_\odot^5}{3 \cdot 5} = - \frac{3}{5} \frac{GM_\odot^2}{R_\odot}$$

0.2

A dimensional analysis with a final answer of $\frac{GM_\odot^2}{R_\odot}$

(0.1)

A final answer of $+\frac{3}{5} \frac{GM_\odot^2}{R_\odot}$ will earn 0.1 mark since the sign has a crucial physical significance.

(0.1)

- (A10) Estimate the maximum possible time τ_{KH} (in years), for which the Sun could have been shining, according to the KH hypothesis. Assume that the luminosity of the Sun has been constant throughout this period. [0.5]

Solution:

$$\tau_{\text{KH}} = \frac{-\Omega}{L_{\odot}}$$

$$\tau_{\text{KH}} = \frac{3GM_{\odot}^2}{5R_{\odot}L_{\odot}} = 1.88 \times 10^7 \text{ years}$$

0.3

0.2


In final answer is negative, deduct 0.2 here.

If there was a sign error in A9, but the answer here finally is positive, no marks deducted in A10.

The τ_{KH} calculated above does not match the age of the solar system estimated from studies of meteorites. This shows that the energy source of the Sun cannot be purely gravitational.

B. Neutrinos from the Sun:

In 1938, Hans Bethe proposed that nuclear fusion of hydrogen into helium in the core of the Sun is the source of its energy. The net nuclear reaction is:



The “electron neutrinos”, ν_e , produced in this reaction may be taken to be massless. They escape the Sun and their detection on Earth confirms the occurrence of nuclear reactions inside the Sun. Energy carried away by the neutrinos can be neglected in this problem.

- (B1) Calculate the flux density, Φ_{ν} , of the number of neutrinos arriving at the Earth, in units of $\text{m}^{-2}\text{s}^{-1}$. The energy released in the above reaction is $\Delta E = 4.0 \times 10^{-12}$ J. Assume that the energy radiated by the Sun is almost entirely due to this reaction. [0.6]

Solution:

$$4 \times 10^{-12} \text{ J} \leftrightarrow 2\nu$$

$$\Rightarrow \Phi_{\nu} = \frac{L_{\odot}}{4\pi d_{\odot}^2 \Delta E} \times 2 = \frac{3.85 \times 10^{26}}{4\pi \times (1.50 \times 10^{11})^2 \times 4.0 \times 10^{-12}} \times 2 = 6.8 \times 10^{14} \text{ m}^{-2}\text{s}^{-1}.$$

0.3

0.3

0.1 for each step, 6.5–7.0 range OK

Travelling from the core of the Sun to the Earth, some of the electron neutrinos, ν_e , are converted to other types of neutrinos, ν_x . The efficiency of the detector for detecting ν_x is 1/6th of its efficiency for detecting ν_e . If there is no neutrino conversion, we expect to detect an average of N_1 neutrinos in a year. However, due to the conversion, an average of N_2 neutrinos (ν_e and ν_x combined) are actually detected per year.

- (B2) In terms of N_1 and N_2 , calculate what fraction, f , of ν_e is converted to ν_x . [0.4]

Solution:

$$\begin{aligned}
N_1 &= \epsilon N_0 \\
N_e &= \epsilon N_0(1-f) \\
N_x &= \epsilon N_0 f/6 \\
N_2 &= N_e + N_x
\end{aligned}$$

OR

$$\begin{aligned}
(1-f)N_1 + \frac{f}{6}N_1 &= N_2 \\
\Rightarrow f &= \frac{6}{5} \left(1 - \frac{N_2}{N_1} \right)
\end{aligned}$$

0.3

0.1

In order to detect neutrinos, large detectors filled with water are constructed. Although the interactions of neutrinos with matter are very rare, occasionally they knock out electrons from water molecules in the detector. These energetic electrons move through water at high speeds, emitting electromagnetic radiation in the process. As long as the speed of such an electron is greater than the speed of light in water (refractive index, n), this radiation, called Cherenkov radiation, is emitted in the shape of a cone.

- (B3) Assume that an electron knocked out by a neutrino loses energy at a constant rate of α per unit time, while it travels through water. If this electron emits Cherenkov radiation for a time Δt , determine the energy imparted to this electron (E_{imparted}) by the neutrino, in terms of $\alpha, \Delta t, n, m_e, c$. (Assume the electron to be at rest before its interaction with the neutrino.)

[2.0]

Solution:

When the electron stops emitting Cherenkov radiation, its speed has reduced to $v_{\text{stop}} = c/n$.

0.6

Its total energy at this time is

$$E_{\text{stop}} = \frac{m_e c^2}{\sqrt{1 - v_{\text{stop}}^2/c^2}} = \frac{nm_e c^2}{\sqrt{n^2 - 1}}$$

0.6

0.3 marks here if the answer is incorrect, but relativistic nature of the problem is understood.

The energy of the electron when it was knocked out is

$$E_{\text{start}} = \alpha \Delta t + \frac{nm_e c^2}{\sqrt{n^2 - 1}}$$

0.4

Before interacting, the energy of the electron was equal to $m_e c^2$.

0.4

Thus, the energy imparted by the neutrino is

$$E_{\text{imparted}} = E_{\text{start}} - m_e c^2 = \alpha \Delta t + \left(\frac{n}{\sqrt{n^2 - 1}} - 1 \right) m_e c^2$$

For partial credit, check which of the above steps have been completed.

The fusion of H into He inside the Sun takes place in several steps. Nucleus of ${}^7\text{Be}$ (rest mass, m_{Be}) is produced in one of these intermediate steps. Subsequently, it can absorb an electron, producing a ${}^7\text{Li}$ nucleus (rest mass $m_{\text{Li}} < m_{\text{Be}}$) and emitting a ν_e . The corresponding nuclear reaction is:



When a Be nucleus ($m_{\text{Be}} = 11.65 \times 10^{-27}$ kg) is at rest and absorbs an electron also at rest, the emitted neutrino has energy $E_\nu = 1.44 \times 10^{-13}$ J. However, the Be nuclei are in random thermal motion due to the temperature T_c at the core of the Sun, and act as moving neutrino sources. As a result, the energy of emitted neutrinos fluctuates with a root mean square value ΔE_{rms} .

- (B4) If $\Delta E_{\text{rms}} = 5.54 \times 10^{-17}$ J, calculate the rms speed of the Be nuclei, V_{Be} and hence estimate T_c .
(Hint: ΔE_{rms} depends on the rms value of the component of velocity along the line of sight.)

[2.0]

Solution:

Moving ${}^7\text{Be}$ nuclei give rise to Doppler effect for neutrinos. Since the fractional change in energy ($\Delta E_{\text{rms}}/E_\nu \sim 10^{-4}$) is small, the Doppler shift may be considered in the non-relativistic limit (a relativistic treatment gives almost same answer). Taking the line of sight along the z -direction,

$$\begin{aligned} \frac{\Delta E_{\text{rms}}}{E_\nu} &= \frac{v_{z,\text{rms}}}{c} && 0.4 \\ &= 3.85 \times 10^{-4} && 0.1 \\ &= \frac{1}{\sqrt{3}} \frac{V_{\text{Be}}}{c} && 0.5 \end{aligned}$$

$$\Rightarrow V_{\text{Be}} = \sqrt{3} \times 3.85 \times 10^{-4} \times 3.00 \times 10^8 \text{ m s}^{-1} = 2.01 \times 10^5 \text{ m s}^{-1}. \quad 0.2$$

The average temperature is obtained by equating the average kinetic energy to the thermal energy.

$$\begin{aligned} \frac{1}{2} m_{\text{Be}} V_{\text{Be}}^2 &= \frac{3}{2} k_B T_c && 0.6 \\ \Rightarrow T_c &= 1.13 \times 10^7 \text{ K} && 0.2 \end{aligned}$$

The prefactor of $k_B T_c$ may be anywhere between 1/2 and 3/2 to get full credit.